

PRACTICE PARAMETERS
CHILD and ADOLESCENT INPATIENT PSYCHIATRIC TREATMENT
A Publication of the Association of Child and Adolescent Psychiatric Nurses
A Division of International Society of Psychiatric Nurses (ISPN)

Introduction

Child and adolescent inpatient psychiatry is a subspecialty in nursing requiring a definable set of skills and abilities. In the inpatient setting, practice centers on the provision of a safe, structured, and supportive environment for seriously ill young people. Their needs are diverse and complex. Care to this vulnerable population is provided by a multidisciplinary team. These practice statements are meant to be broad parameters for defining performance expectations in various aspects of inpatient treatment.

This document describes parameters on assessment, milieu management, cultural considerations, discharge planning, care of special populations (e.g. suicidal, aggressive, maltreated youth) and guidelines on specific practices such as use of seclusion and restraint. These parameters were developed from clinical expertise and a review of the literature and are offered as a supplement to the American Nurses Association (ANA), *Scope and Standards of Psychiatric-Mental Health Practice* (American Nurses Association, 2000). The ANA document serves as the primary guide for clinical nursing practice with these child and adolescent inpatient parameters offering more specific guidance for child and adolescent subspecialty practitioners. This document is only a snapshot of knowledge to date, and practitioners are urged to use their research interpretation skills to critique and incorporate new knowledge into their practice throughout their professional lives.

We appreciate the work of ACAPN members in revising the existing practice parameters first written in 1995: Pamela Galehouse, Chris Hacker, Ellen C. Rindner, Judith Hirsh, Lois Powell, Edilma Yearwood. Thanks to Judith Coucouvanis, Edilma Yearwood and Elizabeth H. Erwin who assisted in the editing of the final product.

Kathleen Delaney, RN, DNSc
Division Director, Association of Child and Adolescent Psychiatric Nurses
A division of ISPN

American Nurses Association (2007). *Scope and Standards of Psychiatric- Mental Health Nursing Practice*. Washington, D. C.: Author.

These guidelines summarize data to inform nurses and mental health staff of the care of children and adolescents during inpatient psychiatric treatment. These guidelines are not intended to serve as the absolute standards of child/adolescent inpatient psychiatric care. Standards are subject to change as our evidence base grows. Adherence to these guidelines should not be interpreted as including all proper methods of care or excluding other acceptable methods.

ACAPN INPATIENT PRACTICE PARAMETERS**CONTENTS**

ASSESSMENT OF THE CHILD AND ADOLESCENT	3
PRINCIPLES OF THERAPEUTIC MILIEU	8
USE OF PHYSICAL RESTRAINTS AND SECLUSION	15
MANAGEMENT OF AGGRESSION	25
ASSESSMENT OF SUICIDE RISK	33
SEXUAL ABUSE	39
TRANSCULTURAL CARE	46
DISCHARGE PLANNING	51

**ASSESSMENT OF THE CHILD AND ADOLESCENT DURING BRIEF
INPATIENT PSYCHIATRIC TREATMENT**

DESCRIPTION/OVERVIEW:

During inpatient hospitalization, psychiatric nurses are afforded a unique assessment opportunity by virtue of their 24-hour presence on the unit. They observe and interact with patients and significant others with the purpose of assessing behaviors, symptoms and responses to intervention in the milieu. Because of their continuous presence nurses have the opportunity to recognize patterns of behaviors, symptoms, deficits, thoughts and beliefs; and to communicate these to other members on the team. Inpatient units typically admit children with a variety of mood, behavioral and neurobiological disorders as well as children with significant physical or mental co-morbidities. Thus assessment must be holistic, and the data collected using reliable and valid methods.

DESIRED OUTCOME:

Assessment data will be compiled that reflects the child's current status, accurately determines symptoms, and describes how the child/adolescent functions in activities of daily living and in therapeutically designed activities. These activities will vary in structure and task demands. Nurses will assess for any difficulties with regulation of affect or behavior associated with situations such as transitions, activities of daily living (e.g. concentrating on homework), visiting with family members, peer interaction, or periods when the pace and noise level of the milieu increases or decreases. Responses to treatment are documented. The nurse also assesses how the child/adolescent views their illnesses and how effective adults are, as they interact, nurture, discipline and support them in managing their acute illness. Finally, assessment will include any potential barriers to discharge to a less restrictive environment and what services or referrals can increase the likelihood of a successful transition.

A. ASSESSMENT

Principle of Care	Practice
<p>The child/adolescent and family can expect: A thorough assessment of current behavior, centered on issues that prompted admission.</p> <p>An opportunity to explain their perceptions of the behaviors and symptoms that prompted admission.</p> <p>A holistic assessment is conducted including physical and psychiatric issues.</p> <p>Any unstable medical conditions are monitored.</p>	<p>The nurse will:</p> <p>Be proficient in understanding developmental, cultural, spirituality and gender differences in the presenting patients.</p> <p>Utilize reliable and valid methods to collect data; synthesize findings and communicate these with other members of the team.</p> <p>Complete a review of systems as specified on the nursing admission data base.</p>

<p>The assessment reflects not only health behaviors or symptoms, but other factors integral to the child's functioning, such as strengths, affect regulation, support system, coping and stress response, information processing, relatedness and control.</p> <p>Assessment includes evaluation of the child's functioning within peer groups, the family, school, and the neighborhood/ community.</p> <p>There will be continuous assessment of any behaviors/thoughts that indicate a danger to self or others. When known risk factors are present anticipatory planning will be done.</p> <p>The plan of care that is developed is based on the assessment and reflects a synthesis of all relevant data.</p> <p>The plan of care reflects patient-centered goals and family involvement in establishing those goals.</p> <p>The patient, family and legal guardians will understand the unit policies about how assessment data is collected and will participate in the process.</p> <p>The assessment process will include a discussion of confidentiality, trust and the nature of the nurse-patient- guardian relationship.</p>	<p>Assess the child/adolescent for:</p> <ul style="list-style-type: none"> -Current behavior, thoughts, mood and affect, and how they compare with parent/child report pre-admission -Review of medications: current, past years, side/adverse effects, dosage adjustments and response -Situational or relational variables that are related to increased or diminished intensity of problem behaviors -Any verbal or non-verbal signs of danger to self or others (See additional guidelines detailed in section on monitoring suicidal behavior) -Substance use: type, amount, frequency, context. Include legal, illegal, over the counter, and psychoactive substances -Legal involvement -Traumatic events (recent and remote) -Achievement of developmental milestones -Functional limitations that may impact discharge planning -Any co-occurring medical conditions. <p>Obtain history of psychiatric treatment.</p> <p>Assess the family for ability to provide for the patient's needs post discharge.</p> <p>Assess for any potential barriers for discharge to the least restrictive environment.</p>
---	---

B. DIAGNOSES

Assessment could result in any number of nursing diagnoses, including but not limited to:

Impaired social interaction

Altered role performance

Altered growth and development

Risk for injury/Risk for violence self directed or other

Self-care deficit

Ineffective coping

C. PLANNING

Principle of Care	Practice
<p>A complete assessment of the patient's behavior requires planning for the on-going assessment of milieu behavior. This milieu assessment of behavior demands:</p> <p>Observing and interacting with the child/adolescent in a variety of milieu settings that includes different levels of stimulation and task demands.</p> <p>Obtaining parent/guardian perceptions of milieu behavior.</p> <p>Scheduling times to discuss staff observations of milieu behaviors with the parent/guardian.</p> <p>Assuring that sufficient numbers of appropriately trained staff are assigned to detect variations in behaviors, affects, functioning and response to treatment.</p> <p>Providing continuity of staff to promote trust and rapport and enable ongoing assessment.</p> <p>Inclusion of knowledge deficits that will inform both treatment and discharge planning.</p>	<p>The nurse will:</p> <p>Plan opportunities to assess functioning in a variety of situations (e.g. with peers, doing structured and unstructured tasks) and determine if variation exists.</p> <p>Include a plan for systematically engaging with primary guardians.</p> <p>Allow multiple staff to assess behavior in a variety of situations.</p> <p>Plan staffing to ensure consistency with patient assignments, facilitate the development of the therapeutic relationship and thus enable assessment of changes in response to treatment.</p> <p>Consult with the literature when necessary to ensure that all relevant factors are assessed.</p> <p>Communicate and collaborate in developing this plan with the multidisciplinary team.</p> <p>Integrate assessment findings and prioritize.</p>

D. IMPLEMENTATION

Principle of Care	Practice
<p>Implementing the assessment process requires the nurse has knowledge of all of the psychiatric disorders, and of the physiological illnesses that have associated psychiatric symptoms.</p>	<p>The nurse will:</p> <p>Identify patterns of behaviors over time and identify antecedents or consequences of behavior.</p> <p>Utilize assessment tools as appropriate to gather quality data.</p>

<p>In assessing youth the nurse acknowledges:</p> <p>The number of stressors prior to admission is related to child's presentations and adjustment in the hospital.</p> <p>Assessments will be individualized to the needs of the child, and will be conducted using therapeutic tools, reliable and valid methods.</p> <p>Therapeutic interactions for the purpose of assessment will take into consideration the nature of the stressors and symptoms, as well as gender or cultural issues that may impact disclosure.</p>	<p>Observe the patient in a variety of milieu and group situations.</p> <p>Discuss incidents and behaviors with patient to ascertain their perceptions and attributions.</p> <p>Discuss observations with parent or guardian to compare and contrast milieu behaviors with home behaviors.</p> <p>Provide constant supervision so that behaviors indicative of danger to self or others can be observed. (See additional guidelines on suicidal patients)</p> <p>Consider the relative impact of stressors in school, peer group and neighborhood on adjustment.</p>
---	---

E. EVALUATION

Principle of Care	Practice
<p>Assessment of behaviors is an ongoing process that provides information useful to diagnosis, treatment, and discharge planning.</p> <p>Inpatient facilities that experience optimal outcomes use a systematic process of risk management, independent auditing of programs and policies, and evaluating their practices against 'best evidence' in the literature.</p>	<p>The nurse will:</p> <p>Communicate staff's assessment of patient's milieu behavior to the treatment team.</p> <p>Conduct and participate in discussions among nursing staff to reach common understanding of the child or adolescent.</p> <p>Refine assessment as new data emerges and/or behaviors or symptoms change.</p> <p>Consult the literature as needed.</p>

RESOURCES

- American Academy of Child and Adolescent Psychiatry (1997). Practice parameters for the psychiatric assessment of children and adolescents. *Journal of the American Academy of Child and Adolescent Psychiatry*, 36 (10 Supplement), 4S-20S.
- Delaney, K.R. (in press). Special Edition: Milieu Assessment of Behavior. *Journal of Child and Adolescent Psychiatric Nursing*.
- Greenspan, S. I. & Greenspan, N. T. (2003). *The clinical interview of the child*. Washington, D. C: American Psychiatric Association.
- Joint Commission on Accreditation of Health Care Organizations (2003). *Assessing cognitive and emotional functioning across the continuum of care*. Oakbrook Terrace, Illinois: Author.
- Kronenberger, W. C., Carter, B. D., & Thomas, D. (1997). Assessment of behavior problems in pediatric inpatient settings: Development of a Pediatric Inpatient Behavior Scale. *Children's Health Care*, 26, 221-232.
- Lyall, D., Hawley, C., & Scott, K. (2004). Nurses' observation scale for inpatient evaluation: Reliability update. *Journal of Advanced Nursing*, 46, 390-394.
- Thienemann, M. (2004). Introducing a structured interview into a clinical setting. *Journal of the American Academy of Child and Adolescent Psychiatry*, 43, 1057-1060.

PRINCIPLES OF THE THERAPEUTIC MILIEU APPLIED TO PSYCHIATRIC HOSPITALIZATION FOR CHILDREN AND ADOLESCENTS

DESCRIPTION/OVERVIEW:

Inpatient psychiatric nurses are the “gatekeepers of the milieu.” They assume responsibility for the milieu’s tone, pace and activity level. They assure that the milieu is safe, structured and supportive. Further, they assure the milieu is engineered so that the child/adolescent succeeds and begins to build a sense of control in their lives. A therapeutic milieu consists of five components: safety, structure, support, involvement and validation. Inpatient units are typically locked and the young person’s movements about the space are restricted (e.g. visitation between patients in their rooms). Containment limits a child/adolescent’s autonomy and freedom of will and separates them from sources of support such as parents and peers. Also, adults possess control that can be intimidating or anxiety provoking for many young patients, especially those who have been victimized. For many youth, the issue of containment is extremely stressful and impacts their coping and behavior on the unit.

Structure in the milieu comes from unit policies and procedures, the practice behavior of the staff, and from incorporating existing knowledge about therapeutic activities into nursing practice. It is based on meeting the developmental needs of the patients. Increasingly structure is informed by principles of family-centered care. The structure provides the backdrop for the nurse to assess responses to hospitalization.

Support, involvement and validation comes from adult staff as well as peers, other participants in unit activities (e.g. parents in a family group) and other visitors to the physical space (e.g. housekeeping staff). Children and adolescents have a right to feel safe and respected, have personal boundaries protected, be reliably and consistently responded to and experience trust in the milieu staff. Staff will collaborate to provide a balance between support, structure and containment.

DESIRED OUTCOME:

Each child or adolescent’s mental and physical health is fostered during their inpatient stay. Through participating in the structured activities of the milieu, the child or adolescent learns coping skills and experiences being efficacious at self-regulation. They will identify unhealthy thoughts and behaviors as they occur and, with the assistance of the nursing staff, modify these behaviors and/or replace them with effective coping skills. The child or adolescent uses peer, parental and staff feedback and teaching from staff to modify maladaptive behaviors. These activities, which often mimic typical activities of daily living, present opportunities for the young person to demonstrate their progress in areas such as impulse control. When participating in milieu treatment, the nurse intervenes in problem areas identified in the plan of care. Nursing staff will build a therapeutic alliance with patients and their families; and serve as a patient advocate. It is important for the environment to be carefully structured to meet both the needs of the individual and the needs of the group. Often, children who are hospitalized have been traumatized. Their emotion regulation and stress responses are often impaired. A critical outcome for these children is that the milieu does not precipitate re-traumatization.

A. ASSESSMENT

Principle of Care	Practice
<p>The milieu will be constructed to meet the requirements for safety, structure, support, involvement, and validation.</p> <p>Young people who have been maltreated or are victims of violence are at high risk for acute stress disorder and require a therapeutic milieu for healing and safety.</p> <p>The child/adolescent and family will experience respect and have his/her needs for safety, personal boundaries and trust assessed and met from the point of admission through discharge.</p> <p>The child/adolescent and family can expect the nursing staff to assess and control the milieu's pace, tone and activity levels from the point of admission through the point of discharge.</p> <p>The structure of the milieu will be targeted to the developmental, physical and psychological needs of the youth who are admitted.</p> <p>Individual expectations will be kept in line with the child's ability to reason, perform and regulate.</p> <p>The unit lives by sensible rules that elicit the child's cooperation and, in as many instances as possible, afford the child choice.</p> <p>Milieu guidelines include includes clear rules and consequences for antisocial actions.</p>	<p>The nurse will: Continuously assess patients' safety needs.</p> <p>Continuously monitor the milieu's tone, pace, activity level.</p> <p>Provide a schedule and list of activities for children and review it with them.</p> <p>Seek patient input on activities and unit guidelines.</p> <p>Assess and respond to boundary conflicts between milieu and therapy staff.</p> <p>Collaborate with the patient and family on patient's needs and how the milieu might address them.</p> <p>Conduct assessments that consider both the needs of the individual young person and the needs of the milieu.</p> <p>Be mindful of any child's history of violence or victimization and its elaboration in milieu behaviors.</p> <p>Detect and prevent periods of group contagion and milieu escalation.</p> <p>Assess the patient group needs and staffing numbers to assure sufficient staff is available to respond to and address youth's needs.</p> <p>Assess the needs of the individual youth against milieu expectations.</p>

B. DIAGNOSIS

Assessment could result in any number of nursing diagnoses, including but not limited to:

High risk for violence, self or other-directed
 Ineffective individual coping related to child abuse or neglect
 Self-esteem disturbance related to unmet dependency needs
 Impaired social interaction related to negative role modeling
 Defensive coping related to low self-esteem
 High level of anxiety related to treatment in a setting other than home or school

B. PLANNING

Principle of Care	Practice
<p>The child/adolescent and family can expect: Members of the treatment team will maintain the safety of the milieu and promote success in coping and functioning with milieu expectations.</p> <p>Planning will be done in the five areas of milieu: safety, structure, support, involvement and validation.</p> <p>The nurse will assess the milieu regularly and intervene to maintain safety and structure, and engage in anticipatory problem solving with other staff to avoid crises.</p> <p>Planning will assure integration of family-centered goals.</p>	<p>The nurse will:</p> <p>Collaborate with the child/adolescent family and treatment team to determine the proposed interventions/strategies for treating the client's basic identified needs, i.e., safety needs and containment of behaviors that are dangerous to self or others.</p> <p>Collaborate with unit management to assess milieu acuity, determine therapeutic activities, and adjust staff levels to assure a safe, structured and supportive milieu, and one where patients succeed in meeting the demands of hospitalization.</p> <p>Adjust staffing numbers to assure sufficient staff is available to respond to and address patient's needs.</p> <p>Assure clinical structures and resources are in place such that all milieu functions (safety, structure, support, involvement, validation) can be consistently enacted.</p>

C. IMPLEMENTATION

Principle of Care	Practice
<p>Youngsters who are hospitalized because they posed a serious danger to self or others require a safe, structured environment which supports their regulation of emotions and behavior.</p> <p>Hospitalized youth who have experienced trauma or maltreatment may have additional problems with arousal patterns, stress response, sleep/rest cycles, memory and attachment. They heal and learn best in an environment of sensible rules where respect and shared decision making support autonomy and reduce stress.</p> <p>In staff's efforts to create order and structure, the child learns that staff is dependable, responsive and predictable.</p> <p>Many hospitalized youngster have social cognitive skill deficits, personal space and boundary limitations, and react strongly to peer rejection, or other interpersonal stressors. The milieu is structured such that staff step in and bolster faltering function so the child succeeds.</p> <p>The youngsters can expect that they will receive positive reinforcement when they behave in a safe manner, follow expectations and participate in treatment activities.</p> <p>The child/adolescent/family can expect to interact with nurses in the milieu and receive on-going, individualized client and family education related to diagnosis, treatment approaches, and psychotropic medications.</p>	<p>The nurse will:</p> <p>Establish a trusting relationship with the child/adolescent/family in a way that conveys interest, respect, and support.</p> <p>Inform the child/adolescent and family of the unit structure, rules and expectations during the course of treatment.</p> <p>Provide a balance of structure and unstructured or quiet time.</p> <p>Strategize with others to maintain a calm, organized milieu.</p> <p>Maintain a positive tone to the environment.</p> <p>Continuously assess the milieu for safety by performing random and routine checks for safety.</p> <p>Maintain staff visibility at a maximum level.</p> <p>Maintain awareness of the child's particular deficits and needs and reliably step in and provide support to the child's regulation or cognitive functions.</p> <p>Reinforce positive/adaptive changes in the client's coping and attempt to increase positive interactions with the child/adolescent.</p> <p>Use milieu interactions and developmentally appropriate teaching tools to provide the child and family education on interventions, diagnoses, and medications.</p>

E. EVALUATION

Principle of Care	Practice
<p>Child/adolescent/family can expect ongoing, timely evaluations of their responses to expressed safety, personal boundary, and trust needs from point of admission to unit through the discharge phase of treatment.</p> <p>Outcomes are evaluated by patient, family and staff to increase satisfaction with services.</p> <p>Staff continually monitor the safety of the milieu and examine any lapses in safety that occur.</p> <p>Staff monitor their response to milieu situations and evaluate the individual's or group's response to the approach.</p>	<p>The nurse will</p> <p>Monitor/evaluate the client/family's response to treatment on an ongoing basis and document this evaluation of responses.</p> <p>Review (at shift report) which milieu management strategies have been effective and which have not and revise strategies appropriately.</p> <p>Engage in anticipatory problem solving (e.g. separating peers with a history of conflict) to increase success in activities.</p> <p>Monitor and evaluate responses to treatment on an ongoing basis and share evaluation with multidisciplinary team members.</p> <p>Review and integrate patient/family perceptions of care including discharge satisfaction data.</p> <p>Collaboratively identify client/family/community support systems following discharge to promote the continuation of behavioral change.</p>

RESOURCES

- Bloom, S. L., Bennington-Davis, M., Farragher, B., McCorkle, D., Nice-Martini, K., & Wellbank, K. (2003). Multiple opportunities for creating sanctuary. *Psychiatric Quarterly*, 74, 173-190.
- Causey, D, McKay, M, Rosenthal, C & Darnell, C. (1998). Assessment of hospital related stress in children and adolescents admitted to a psychiatric unit. *Journal of Child and Adolescent Psychiatric and Mental Health Nursing*, 11(3), 135-145.
- Cummings, M., Waller, D. Johnson, C., Bradley, K., Leatherwood, D. & Guzetta, C. (2001). Developing and implementing a comprehensive program for children and adolescents with eating disorders. *Journal of Child and Adolescent Psychiatric and Mental Health Nursing*, 14, 167-177.
- Delaney, K. R. (1994). Calming an escalated inpatient milieu. *Journal of Child and Adolescent Psychiatric and Mental Health Nursing*, 7(3), 5-13.
- Delaney, K. R. (1997). Milieu therapy: A therapeutic loophole. *Perspectives in Psychiatric Nursing*, 33(2), 19-28.
- Delaney, K.R., Perraud, S., & Pitula, C. (2000). Brief psychiatric hospitalization and process description: What will nursing add? *Journal of Child and Adolescent Psychosocial Nursing*, 3, 7-13.
- Delaney, K. (2002). Point of view. Inpatient psychiatric nursing: set up to stagnate? *Journal of the American Psychiatric Nurses Association* 8(4), 130-134.
- Erwin, E.H. (2002). Perceptions of relevant social problems. *Journal of Child and Adolescent Psychiatric and Mental Health Nursing*, 15(1), 24-34.
- Geanellos, R. (2000). The milieu and milieu therapy in adolescent mental health nursing. *International Journal of Psychiatric Nursing Research*, 5(3), 638-48.
- Lawson, L. (1998). Milieu management of traumatized youngsters. *Journal of Child and Adolescent Psychiatric and Mental Health Nursing*, 11(3), 99-106.
- Marriage, K., Petrie, P., & Worling, D.(2001). Consumer satisfaction with an adolescent inpatient psychiatric unit. *Canadian Journal of Psychiatry*, 46, 969-975.
- Mohr, W. K. (2003). Johnson's Psychiatric Mental-Health Nursing (5th ed.). Philadelphia: Lippincott William & Wilkins.

- Regan, K. M., Curtin, C., & Vorderer, L. (2006). Paradigm shifts in inpatient psychiatric care of children: Approaching child-and-family centered care. *Journal of Child and Adolescent Psychiatric Nursing, 19*, 29-40.
- Rivard, J. C., McCorkle, D., Duncan, M. E., Pasquale, L. E., Bloom, S. L., & Abramovitz, R. (2004). Implementing a trauma recovery model for youths in residential treatment. *Child and Adolescent Social Work Journal, 21*, 529-550.
- Romer, D. (2003). *Reducing adolescent risk*. Thousand Oaks: Sage Publishers.
- Soth, N. B. (1997). *Informed treatment: Milieu management in psychiatric hospitals and Residential treatment centers*. Blue Ridge Summit, PA: Scarecrow Press.
- Sharer, K. (1999). Nurse-parent relationship building in child psychiatric units. *Journal of Child and Adolescent Psychosocial Nursing, 4*, 153-167.
- Townsend, M. C. (1994). *Nursing diagnoses in psychiatric nursing: A pocket guide for Care plan construction*. Philadelphia: F.A. Davis Company.
- Townsend, M. C. (1996). Interactive therapies and methods of implementation. In K. Fortinash & P. Holoday-Worret (Eds.), *Psychiatric-Mental Health Nursing* (pp. 502-530). St. Louis: Mosby-Year-Book, Inc

GUIDELINES FOR THE USE OF RESTRAINT AND SECLUSION DURING CHILD AND ADOLESCENT INPATIENT TREATMENT

DESCRIPTION/OVERVIEW:

Restraint means restricting the freedom of movement or normal access of a person to his/her own body by means of a manual method or physical or mechanical device (Health Care Financing Administration [HCFA], 1999). Seclusion is the involuntary confinement of a patient to either a room or area such that the person is physically prevented from leaving (HCFA, 1999). Potential and actual dangerous behavior requires immediate evaluation and intervention to prevent and manage harm to self and/others. Restraint and seclusion have not been found to be useful as interventions for modifying behavior. However, the literature does support the use of restraint and seclusion as safety interventions. Restraint and seclusion shall be non-punitive and non-coercive and used only in situations that pose a significant danger to the patient or others. Types of restraint include: 1). Mechanical restraints-body movements are contained, restricted or restrained by use of a mechanical device; 2). Physical restraints - body movements are contained or restricted or restrained by the physical actions of a staff person. 3). Chemical restraints-use of a drug or medication that to control behavior or restrict the individual's freedom of movement that is not a standard treatment for the individual's medical or psychiatric condition.

Following restraint/seclusion incidents it is important to process the event with the patient to understand how it is perceived by him/her; and to determine what other interventions could have been attempted and thus avoid restraint in the future. A comprehensive interdisciplinary team approach should be utilized to identify alternate interventions, reduce restraint/seclusion and ensure safety of clients. Restraint and seclusion must be utilized within the guidelines or policies of each agency and also meet the requirements of accrediting and licensing organizations.

Guidelines and standards of practice concerning the use of seclusion and restraint during inpatient hospitalization have been created by Center for Medicaid and Medicare Services (CMS) (formerly HCFA), State Mental Health Acts, and professional organizations such as the International Society for Psychiatric Mental Health Nurses and the American Psychiatric Nurses Association. When applicable, child inpatient nurses should be aware of and comply with the content of these regulations. The guidelines presented here are intended as a supplement to these more extensive parameters and regulations.

DESIRED OUTCOME:

The client will cease injurious behavior and be able to maintain control without the use of restraint or seclusion. The client will learn alternative ways to respond to intense emotions such as anger or frustration; and process experiences that prompt emotional arousal. The staff will be successful in assessing potential situations that may result in restraint or seclusion and receive ongoing training and supervision around the use of restrictive measures and alternative approaches.

A. ASSESSMENT

Principle of Care	Practice
<p>The child/adolescent and family can expect that:</p> <p>A thorough assessment of the child/adolescent's ability to control his/her behavior and motivation for harming self and others will be done prior to the decision to use restraints or seclusion.</p> <p>The welfare and safety of the child/adolescent will be the prime consideration.</p> <p>Nursing staff will assess any situations that may trigger aggression and act to interrupt escalation.</p> <p>Feedback on effective de-escalation methods will be collected during the assessment period.</p>	<p>The nurse will assess the child or adolescent for:</p> <p>Current and historical potential for violence to self or others.</p> <p>Response to less restrictive interventions or to medications that have been helpful in the past.</p> <p>De-escalation methods that have been used successfully in the past.</p> <p>Contraindications or special considerations in using restraint or seclusion.</p> <p>History of trauma, including potential triggers to aggression towards self or others.</p> <p>Any injuries following restraint or seclusion.</p> <p>A clinically privileged nurse will communicate assessment findings to the physician prior to restraining/secluding, when possible, and follow all state, federal and regulatory guidelines concerning physician and LIP notification and responsibilities.</p>

B. DIAGNOSIS

Assessment could result in any number of nursing diagnoses, including but not limited to:

- Potential or high risk for violence toward others
- Potential or high risk for violence toward self
- Potential or high risk for explosive behavior
- Potential or high risk for poor impulse control

C. PLANNING

Principle of Care	Practice
<p>The child/adolescent/family can expect to:</p> <p>Participate collaboratively in planning responses to aggressive behaviors.</p> <p>Be informed of the procedure used to restrain or seclude clients.</p> <p>Be informed, within a reasonable period of time when the child/adolescent is placed in restraints, locked seclusion, or chemical restraint.</p> <p>Staff will be trained on alternative interventions that will decrease levels of patient stress and frustration, and avoid group contagion.</p> <p>Administration will consult with specialists in changing the physical environment to increase safety and monitoring.</p> <p>Administration will review policies and practices to determine any ways to make them less restrictive to staff and patients.</p> <p>Staff will support family involvement in care and in their decisions of how to help their child regulate.</p>	<p>The nurse will:</p> <p>Collaborate with the child/ adolescent/ family by explaining the potential need for restraint or seclusion.</p> <p>Explain behaviors that would result in restraint or seclusion and assess understanding.</p> <p>Describe the restraint/seclusion process and show the patient/family the seclusion room. Ask the family preferences if emergency measures are needed for safety of patient or others.</p> <p>Collaborate with the child/adolescent and family to develop a milieu approach to prevent behavioral and/or affective dyscontrol.</p> <p>Develop a plan with the client to avoid or cope effectively with environmental and internal triggers.</p> <p>While calm, provide options for patients to select as their preferred method of handling lack of control, including time out, verbal interactions with staff, PRN medications, and relaxation techniques.</p> <p>Track behavior to identify any patterns to an individual patient escalation.</p> <p>Assure that all staff involved in restraining or secluding a patient are trained and certified in the use of restraints/seclusion according to agency requirements.</p> <p>As clinically appropriate, follow the family wishes regarding the use of restraints or seclusion.</p>

D. IMPLEMENTATION

Principle of Care	Practice
<p>While in restraints or seclusion, the child/adolescent can expect:</p> <p>To be treated in a safe and respectful manner.</p> <p>To be constantly observed.</p> <p>To receive food, water and toileting opportunities at least every 2 hours.</p> <p>To be evaluated by a licensed independent practitioner within one hour</p> <p>That all hospital, state and regulatory guidelines on restraint procedures will be adhered to.</p> <p>Many alternatives exist to manage behaviors that have been shown to be effective. Nursing staff behavior will consistently utilize these interventions.</p> <p>Restraint and seclusion are never initiated/conducted simultaneously.</p> <p>Institutions will support a ‘restraint free’ environment, including the use of alternative behaviors and consulting with others and the literature on ways to change policy and practice.</p>	<p>The nurse will:</p> <p>Follow the agency’s procedures for restraints and seclusion, as stipulated in JACHO, CMS (formerly HCFA) guidelines, and the State Mental Health Codes.</p> <p>Use less restrictive interventions prior to the use of restraint or seclusion.</p> <p>Give clear expectations to the patient about the expected behavior.</p> <p>Assure that all restraint and seclusion episodes are initiated and terminated in a safe and therapeutic manner by trained staff.</p> <p>Use sufficient numbers of staff with crisis management training to move the child/adolescent into seclusion or restraint.</p> <p>Clearly communicate to the client the behaviors required of them to end the restraint or seclusion episode.</p> <p>Maintain constant observation of the client and document at least every 15 minutes the client’s behavior and response to restraints or seclusion.</p> <p>Remove the client from restraints/seclusion immediately when he/she meets the criteria.</p> <p>When the youth regains control, develop with them a consistent, step-by-step plan for reintegration into milieu activities.</p>

D. EVALUATION

Principle of Care	Practice
<p>The child/adolescent/family can expect to participate in the evaluation of the use of restraints or seclusion.</p> <p>The institution will support an aggression task force to look at practices, staff training and supervision, perceptions, and alternative interventions.</p> <p>Staff should understand that when young people are restrained or secluded there can be negative effects, such as physical injury, psychological distress, and damage to the nurse-patient relationship.</p> <p>Unit management recognizes that instances of restraint and seclusion are also potentially traumatizing for the staff, (physically and emotionally), and for the other patients.</p> <p>Institutional support should exist for research that identifies nurse and patient behavior and beliefs before, during and after restraint/seclusion incidents.</p> <p>The institution will assure a process for on-going quality assurance review of all episodes of restraint or seclusion.</p>	<p>The nurse will: Continuously monitor/evaluate the client's response to restraints or seclusion.</p> <p>Following the restraint or seclusion incident debrief with the client to identify and review behavioral problems, biologic precipitants and environmentally specific triggers that led to restraint.</p> <p>Assess if any injuries occurred during the incident and identify contributing factors.</p> <p>Explore feelings and perceptions with the intent of minimizing potential negative effects of the restraint or seclusion and to restore the nurse-patient relationship.</p> <p>Structure debriefing to facilitate the recognition of earlier symptom recognition and de-escalation; to improve affect regulation, impulse control and motivation for safety; and to promote problem-solving and conflict resolution skills.</p> <p>Debrief with other patients as needed (e.g. witnesses to the event).</p> <p>Participate in quality assurance review of the use of restraint and seclusion by collecting specific data about the clinical endpoints of restraint in order to identify patterns of use.</p> <p>Data collected will include the child/adolescent and family's perception and reaction to restraint or seclusion.</p>

RESOURCES

- Allen, J. (2000). Seclusion and restraint of children: A literature review. *Journal of Child and Adolescent Psychiatric Nursing, 13*, 159-167.
- Alty, A. (1997). Nurses' learning experience and expressed opinions regarding seclusion practice within one NHS trust. *Journal of Advanced Nursing, 25*, 786-793.
- American Academy of Child and Adolescent Psychiatry (AACAP) (2002). Practice parameter for the prevention and management of aggressive behavior in child and adolescent psychiatric institutions with special reference to seclusion and restraint. *Journal of the American Academy of Child and Adolescent Psychiatry, 41*, 4S-25S.
- American Academy of Pediatrics (1997). The use of physical restraint interventions for children and adolescents in the acute care setting. *Pediatrics, 99*, 497-498.
- American Psychiatric Association (APA) (2003). APA achievement awards: 2003 APA Gold award: A more compassionate model for treating children with severe mental illness. *Psychiatric Services, 54*, 1529-1531.
- American Psychiatric Nurses Association (APNA) (2000). *Position statement on the use of seclusion and restraint*. Washington, D.C.: Author.
- Berrios, C. D. & Jacobowitz, W. H. (1998). Therapeutic holding: Outcomes of a pilot study. *Journal of Psychosocial Nursing, 36(8)*, 14-18.
- Bloom, S. L., Bennington-Davis, M., Farragher, B., McCorkle, D., Nice-Martini, K., & Wellbank, K. (2003). Multiple opportunities for creating sanctuary. *Psychiatric Quarterly, 74*, 173-190.
- Bower, F. L., McCullough, C. S., & Timmons, M. E. (2003). A synthesis of what we know about the use of physical restraints and seclusion with patients in psychiatric and acute care settings: 2003 update. *Online Journal of Knowledge Synthesis for Nursing, 10*, document 1.
- Bowers, L., Alexander, J., Simpson, A., Ryan, C., & Carr-Walker, P. (2004) Cultures of psychiatry and the professional socialization process: The case of containment methods for disturbed patients. *Nurse Education Today, 24*, 435-442
- Calabro, K., Mackey, T. A., & Williams, S. (2002). Evaluation of training designed to prevent and manage patient violence. *Issues in Mental Health Nursing, 23*, 3-15.

- Canatsey, K. & Roper, J. M. (1997). Removal from stimuli for crisis intervention: Using least restrictive methods to improve the quality of patient care. *Issues in Mental Health Nursing, 18*, 35-44.
- Cowin, L, Davies, R., Estall, G., Berlin, T., Fitzgerald, M., & Hoot, S. (2003). De-escalating aggression and violence in the mental health setting. *International Journal of Mental Health Nursing, 12*, 64-73.
- Currier, G. W. (2003). The controversy over chemical restraint in acute care psychiatry. *Journal of Psychiatric Practice, 9*, 59-70.
- Delaney, K.R. (2001). Developing a restraint reduction program for child/adolescent inpatient treatment. *Journal of Child and Adolescent Psychiatric Nursing, 14*(3), 128-140.
- Delaney, K. R. & Fogg, L. (2005). Patient characteristics and setting variables related to use of restraint on four inpatient psychiatric units. *Psychiatric Services, 56*, 186-193.
- Delaney, K. R. (2006). Evidenced-Based Practice: Reduction of Restraint and Seclusion Use during Child and Adolescent Psychiatric Inpatient Treatment. *Worldviews on Evidenced Based Practice in Nursing, 5*, 19-30.
- Department of Health and Human Services, Center for Medicare and Medicaid Services (2006). 42 CFR Part 482. Medicare and Medicaid Programs: Hospital conditions of participation: Patients' Rights; final rule. Retrieved on January 1, 2007 from <https://www.cms.hhs.gov/CFCsAndCoPs/downloads/finalpatientrightsrule.pdf>
- Donat, D. (1998). Impact of mandatory behavioral consultation on seclusion/restraint utilization in a psychiatric hospital. *Journal of Behavior Therapy and Experimental Psychology, 29*, 13-19.
- Donat, D. C (2003). An analysis of successful efforts to reduce the use of seclusion and restraint at a public psychiatric hospital. *Psychiatric Services, 54*, 1119-1123.
- Donovan, A., Plant, R., Peller, A., Siegel, L. & Martin, A. (2003). Two-year trends in the use of seclusion and restraint among psychiatrically hospitalized youth. *Psychiatric Services, 54*, 987-993.
- Donovan, A., Siegel, L., Zera, G., Plant, R., & Martin, A. (2003). Child and Adolescent Psychiatry: Seclusion and restraint reform: An initiative by a child and adolescent psychiatric unit. *Psychiatric Services, 54*, 958-959.

- dosReis, S., Barnett, S., Love, R. C., & Riddle, M. A. (2003). A guide for managing acute aggressive behavior of youths in residential and inpatient treatment facilities. *Psychiatric Services, 54*, 1357-1363
- Fryer, M. A., Beech, M., & Byrne, G. J. A. (2004). Seclusion use with children and adolescents: An Australian experience. *Australian and New Zealand Journal of Psychiatry, 38*, 26-33.
- Goren, S., Abraham, I., & Doyle, N. (1996). Reducing violence in a child psychiatric hospital through planned organizational change. *Journal of Child and Adolescent Psychiatric Nursing, 9*(2), 27-37.
- Goren, S. & Curtis, W. (1996). Staff members' beliefs about seclusion and restraint in child psychiatric hospitals. *Journal of Child and Adolescent Psychiatric Nursing, 9*(4), 7-14.
- Goren, S., Singh, N. N., & Best, A. M. (1993). The aggression-coercion cycle: Use of seclusion and restraint in a child psychiatric hospital. *Journal of Child and Family Studies, 2*(1), 61-73.
- Halamandaris, P. & Anderson, T. (1999). Children and adolescents in the psychiatric emergency setting. *Psychiatric Clinics of North America, 22*, 865-874.
- Hamrin, V. (1998). Psychiatric interviews with pediatric gunshot patients. *Journal of Child and Adolescent Psychiatric Nursing, 11*, 61-68.
- Holzworth, R. J. & Willis, C. E. (1999). Nurses' judgments regarding seclusion and restraint of psychiatric patients: A social judgment analysis. *Research in Nursing and Health, 22*, 189-201.
- Huckshorn, K. A. (2004). Reducing seclusion and restraint use in mental health settings: Core strategies for prevention. *Journal of Psychosocial Nursing and Mental Health Services, 42*(9), 22-33.
- International Society for Psychiatric-Mental Health Nursing (ISPN) (1999). *Position on the rights of children in treatment setting*. Retrieved September 1, 2001, from <http://www.ispn-psych.org/docs/99childrens-rights.pdf>
- International Society for Psychiatric-Mental Health Nursing (ISPN) (1999). *Position statement of the use of restraint and seclusion*. Retrieved September 1, 2001, from <http://www.ispn-psych.org/docs/99Restraint-Seclusion.pdf>
- Lawson, L. (1998). Milieu management of traumatized youngsters. *Journal of Child and Adolescent Psychiatric and Mental Health Nursing, 11*(3), 99-106.

- LeBel, J., Stromberg, N., Duckworth, K., Kerzner, J., Goldstein, R., Weeks, M., Harper, G., Sudders, M. (2004). Child and adolescent inpatient restraint reduction: A state initiative to promote strength-based care. *Journal of the American Academy of Child and Adolescent Psychiatry, 43*, 37-45.
- Lee, S., Gray, R., Gournay, K., Wright, S., Parr, A. M., & Sayer, J. (2003). Views of nursing staff on the use of physical restraint. *Journal of Psychiatric and Mental Health Nursing, 10*, 425-430.
- Luiselli, J. K., Bastien, J. S., & Putman, R. F. (1998). Behavioral assessment and analysis of mechanical restraint utilization on a psychiatric, child and adolescent inpatient setting. *Behavioral Interventions, 13*, 147-155.
- Luiselli, J. K., Kane, A., Treml, T. & Young, N. (2000). Behavioral intervention to reduce physical restraint of adolescents with developmental disability. *Behavioral Interventions, 15*, 317-330.
- Mohr, W. K., Mahon, M. M., & Noone, M. J. (1998). A restraint on restraints: The need to reconsider the use of restrictive measures. *Archives of Psychiatric Nursing, 12*, 95-106.
- Mohr, W. K., Petti, T. A., & Mohr, B. D. (2003). Adverse effects associated with physical restraint. *Canadian Journal of Psychiatry, 48*, 330-337.
- National Association of Psychiatric Health Systems (NAPHS)(2003). *Success stories and ideas for reducing restraint/seclusion in behavioral health*. Washington, D. C.: Author.
- National Association of State Mental Health Program Directors (NASMHPD) 1999). *NASMHPD position statement on seclusion and restraint*. Retrieved March 1, 2005, from http://www.nasmhpd.org/general_files/publications/med_directors_pubs/Seclusion_restraint
- Nijman, H., Bowers, L., Oud, N., & Jansen, G. (2005). Psychiatric nurses experiences with inpatient aggression. *Aggressive Behavior, 31*, 217-227.
- Olofsson, B. & Norberg, A. (2001). Experiences of coercion in psychiatric care as narrated by patients, nurses and physicians. *Journal of Advanced Nursing, 33*, 89-97.

- Petti, T. A., Stigler, K. A., Gardner-Haycox, J., & Dumlao, S. (2003). Perceptions of P.R.N. psychotropic medications by hospitalized child and adolescent recipients. *Journal of the American Academy of Child and Adolescent Psychiatry, 42*, 434-441.
- Rutledge, D. N., & Pravikoff, D. S. (2003) Use of restraints. Part 2. Restraints and seclusion in psychiatric institutions. *The On-Line Journal of Clinical Innovations, 6(3)*, 1-56.
- Ryan, E. P., Hart, V. S., Messick, D. L., Aaron, J., & Burnette, M. (2004). A prospective study of assault against staff by youths in a state psychiatric hospital. *Psychiatric Services, 55*, 665-670.
- Singh, N. N., Singh, S. D., Davis, C. M., Latham, L.L., & Ayers, J. G. (1999). Reconsidering the use of seclusion and restraints in inpatient child and adult psychiatry. *Journal of Child and Family Studies, 8*, 243-253.
- Sourander, A., Aurela, A. & Piha, J. (1996). Therapeutic holding in child and adolescent psychiatric inpatient treatment. *Nordic Journal of Psychiatry, 50*, 375-380.
- Sourander, A., Ellila, H., Valimaki, M., & Piha, J. (2002). Use of holding, restraints, seclusion and time-out in child and adolescent psychiatric in-patient treatment. *European Journal of Child and Adolescent Psychiatry, 11*, 162-167.
- Sourander, A., Helenius, H., & Piha, J. (1995). Treatment difficulty and physical restraint in a child psychiatric setting. *Psychiatria Fennica, 26*, 198-206.
- Stirling, C. & McHugh, A. (1997). Developing a non-aversive intervention strategy in the management of aggression and violence for people with learning disabilities using natural therapeutic holding. *Journal of Advanced Nursing, 17*, 503-509.
- Taxis, J. C. (2002). Ethics and praxis: Alternative strategies to physical restraint and seclusion in a psychiatric setting. *Issues in Mental Health Nursing, 23*, 157-170.
- Troutman, B., Myers, K., Borchardt, C., Kowalski, R., & Bubrick, J. (1998). Case study: When restraints are the least restrictive alternative for managing aggression. *Journal of the American Academy of Child and Adolescent Psychiatric Inpatients, 37*, 434-444.
- Wynn, R. (2002). Medicate, restraint or seclude? Strategies for dealing with violent and threatening behavior in a Norwegian university psychiatric hospital. *Scandinavian Journal of Caring Science, 16*, 287-291.
- Wynn, R. (2003). Staff's attitudes to the use of restraint and seclusion in a Norwegian university psychiatric hospital. *Nordic Journal of Psychiatry, 57*, 453-459.

GUIDELINES FOR THE MANAGEMENT OF AGGRESSION DURING CHILD AND ADOLESCENT INPATIENT PSYCHIATRIC TREATMENT

DESCRIPTION/OVERVIEW:

Aggressive behavior is a common presenting problem with children and adolescents that are referred for psychiatric treatment and is the result of many factors. Among these factors are mental illness; environmental influences; genetic predisposition; and situational variables such as anxiety, fear and affect dysregulation. All acts of aggression are critical communication from children/adolescents. However, it is important to distinguish the motivating factors. There is a difference between reactive aggression, often the result of affect dysregulation or internally driven agitation, and proactive aggression which is often purposive and violent. Interventions for reactive aggression are very different from interventions for proactive aggression. Successful treatment plans to help children/adolescents manage aggressive behavior begin with a thorough assessment of potential and actual aggressive or dangerous behavior at admission. Triggers to escalation are identified as well. It is important to recognize the role of milieu stimulation in escalating situations.

DESIRED OUTCOME:

The child/adolescent will be able to express feelings including negative emotions such as anger and aggression. Moreover they will learn to express these feelings in a socially acceptable manner. During times of anger or aggression the child/adolescent will be free from injury and prevented from harming others. Over the course of hospitalization they will learn self-management /self-soothing techniques. Finally, the child/adolescent will be able to process his/her feelings and perceptions with staff after an episode of aggression.

A. ASSESSMENT

Principle of Care	Practice
At admission and throughout the course of treatment the child/adolescent/family can expect to participate in an assessment of their child's coping responses in various situations, including aggressive behavior and of situations that prompt dyscontrol.	The nurse, in collaboration with the child/adolescent/family will assess the potential for aggression including: -Precipitants that may trigger aggressive behaviors -Type (self, other, property, verbal) -Intensity/Duration/Level (mild, moderate, severe)
Reactive aggression will be managed with a calm milieu, and young people will learn and practice coping strategies to achieve successful affect regulation.	-Proactive versus reactive aggression -Frequency of episodes -Prior successful interventions (verbal, medication, external controls)
Proactive aggression will be managed by the structure and containment of the	-Aftermath of aggression on aggressor, victim and witnesses

<p>milieu and reinforcement of expectations.</p> <p>Staff working with aggressive youth must recognize behaviors in themselves and during interactions with children that precede violent behavior. Recognition of such behaviors is the key to developing effective interventions.</p> <p>Staff should understand the principles of collaborative problem solving.</p>	<p>- Recognition of behaviors that would indicate to staff that patient is getting upset, i.e., pacing, isolating in room.</p> <p>The nurse's assessment of aggressive behavior will consider the developmental level of the child/adolescent.</p> <p>Assessment will include milieu factors that may produce escalating situations.</p> <p>The nurse will assess mental status regularly, monitor the milieu and intervene to maintain safety and structure, and engage in anticipatory problem solving with other staff to avoid aggression.</p>
---	--

B. DIAGNOSIS

Assessment could result in any number of nursing diagnoses, including but not limited to:

Ineffective individual coping
 Disturbed thought processes
 Powerlessness
 Anxiety related to knowledge deficit
 Aggression directed at self or others
 Impaired social interaction skills
 Self-esteem disturbance

C. PLANNING

Principle of Care	Practice
<p>The child/adolescent/family can expect to participate collaboratively in the plan of care regarding episodes of aggression and extreme affect dysregulation.</p> <p>For reactive aggression, a calm milieu with structure, personal space boundaries, decreased stimulation, reassurance and encouragement will be provided.</p> <p>Interactions occurring on the inpatient unit may closely resemble coercive</p>	<p>The nurse will:</p> <p>Collaborate with the child/adolescent/family in developing the treatment goal and intervention strategies for treating aggressive or dysregulated behavior.</p> <p>Collaborate with the patient to develop a realistic plan to prevent and/or decrease the level of aggression.</p> <p>If indicated, include the child/adolescent/family in selecting appropriate</p>

patterns between adults and client, thus requiring staff sensitivity to the patients' reactions to limit setting.	medication for aggressive behavior. Anticipate milieu needs while planning a individual and group therapeutic activities.
---	--

D. IMPLEMENTATION

Principle of Care	Practice
<p>The child/adolescent/family can expect to be treated in a safe, supportive and nonjudgmental manner.</p> <p>The nurse will use the institutions' established methods of assisting children/adolescent with managing their behavior.</p> <p>Milieu stimulation is managed in a proactive and responsive manner.</p> <p>The milieu should be structured to provide choice, eliminate threat and to help children achieve control.</p> <p>Specific skills have been shown to be effective for people with proactive aggression. These include cognitive restructuring, relaxation training, modeling techniques, behavior therapy and collaborative problem solving.</p> <p>Skills are taught during periods of relative calm and openness to material and reinforced in interactions throughout the day.</p> <p>The patient will learn to recognize aggression triggers, discuss typical responses to negative experiences and engage in problem solving to manage stress.</p> <p>In responding to escalating situations staff will intervene utilizing the least,</p>	<p>The nurse will: Assure that all acts of aggression are managed in a safe and therapeutic manner and in accordance with agency policy.</p> <p>Utilize therapeutic communication skills when intervening with an aggressive youth.</p> <p>Take into consideration trauma history and be gender sensitive.</p> <p>Recognize and base interventions upon, the child/adolescents state of control using the least restrictive intervention.</p> <p>Utilize redirection and empathy in the early stages of escalating behavior.</p> <p>Provide positive feedback to the patient for making reasonable choices in an escalating situation.</p> <p>Give the client choices and as broad a role in decision making as clinically appropriate.</p> <p>Recognize and assist patient with utilizing strengths to assist with management of affects, e.g. drawing, playing music.</p> <p>Utilize episodes of affect dysregulation to teach the client to identify and label</p>

<p>restrictive method that has the potential to decrease the tension/negative emotion.</p> <p>All staff should receive training in non-violent de-escalation techniques.</p>	<p>feelings, and more socially acceptable means of expressing anger.</p> <p>Process all aggressive episodes with the client to review precipitants, perceptions, and alternative coping strategies.</p> <p>Assure that adequate and appropriately trained staff are present during aggressive episodes.</p> <p>Proactively control the tone, pace and activity level of the milieu to keep stimulation in an acceptable range.</p>
--	--

E. EVALUATION

Principle of Care	Practice
<p>The child/adolescent/family can expect to participate in the evaluation of progress towards attaining treatment goal(s) relating to the management of aggressive behavior.</p> <p>There is consistent report of improved affect regulation, and no incidents of dangerous behavior to self or others.</p> <p>When stressors are experienced, the patient will utilize learned coping resources and experience behavioral and affective control.</p>	<p>The nurse will monitor, evaluate, and document client response to interventions for management of emotion dysregulation, including aggressive behavior.</p> <p>The team (including staff, patient and family) will revise approaches to management of aggression when indicated.</p>

RESOURCES

- Aldwin, C. A. (1994). *Stress, coping and development: An integrative perspective*. New York: Guilford.
- American Academy of Child and Adolescent Psychiatry(2002). Practice parameter for the prevention and management of aggressive behavior in child and adolescent psychiatric institutions with special reference to seclusion and restraint. *Journal of the American Academy of Child and Adolescent Psychiatry*, 41, 4S–25S.
- Barkley, R. A., (1997). *Defiant children: A clinician's manual for assessment and parent training. Second edition*. New York: Guilford Press.
- Berkowitz, L. (1994). Is something missing? Some observations prompted by the cognitive-neoassociationist view of anger and emotional aggression. In L. R. Huesmann (Ed.) *Aggressive behavior: Current Perspectives* (pp. 35-57). New York: Plenum Press.
- Brendgen, M., Vitaro, F., Tremblay, R. E. & Lavoie, F. (2001). Reactive and proactive aggression: Predictions to physical violence in different contexts and moderating effects of parental monitoring and caregiving behavior. *Journal of Abnormal Child Psychology*, 29, 293-304.
- Causey, D. L., McKay, M., Rosenthal, C. & Darnell, C. (1998). Assessment of hospital related stress in children and adolescents admitted to a psychiatric inpatient unit. *Journal of Child and Adolescent Psychiatric Nursing*, 11, 135-145.
- Carlsson, G., Dahlberg, K., & Drew, N. (2000). Encountering violence and aggression in mental health nursing: A phenomenological study of tacit caring knowledge. *Issues in Mental Health Nursing*, 21, 533-545.
- Conner, D. F., Steingard, R. J., Anderson, J. J., & Melloni, R. H. (2003). Gender differences in reactive and proactive aggression. *Child Psychiatry and Human Development*, 33, 279-294.
- Crick, N. R. & Dodge, K. A. (1996). Social information processing mechanisms in reactive and proactive aggression. *Child Development*, 67, 993-1002.
- Delaney J., Cleary M., Jordan R. & Horsfall J. (2001). An exploratory investigation into the nursing management of aggression in acute psychiatric settings. *Journal of Psychiatric and Mental Health Nursing*, 8, 77–84.
- Delaney K.R. (1994). Calming an escalated milieu. *Journal of Child and Adolescent Psychiatric Nursing*, 7(3), 5–13.
- Dodge, K. A., & Crick, N. R. (1990). Social information-processing bases of aggressive

- behavior in children. *Personality and Social Psychology Bulletin*, 16, 8-22.
- Dodge, K. A., Lochman, J. E., Harnish, J. D., Bates, J. E., & Pettit, G. S. (1997). Reactive and proactive aggression in school children and psychiatrically impaired chronically assaultive youth. *Journal of Abnormal Psychology*, 106, 37-51.
- Duxbury J. (2002). An evaluation of staff and patient views of and strategies employed to manage inpatient aggression and violence on one mental health unit: A pluralistic design. *Journal of Psychiatric and Mental Health Nursing*, 9, 325-337.
- Eisenberg, N., Fabes, R. A., & Guthrie, I. K. (1997). Coping with stress: The roles of regulation and development. In S. A. Wolchik & I. N. Sandler (Eds.), *Handbook of children's coping: Linking theory and intervention* (pp. 41-70). New York: Plenum Press.
- Elias, M. J. & Tobias, S. E. (1996). *Social Problem Solving: Interventions in the schools*. New York: Guilford Press.
- Gunnar, M. R., & Cheatham, C. L. (2003). Brain and behavior interface: Stress and the developing brain. *Infant Mental Health Journal*, 24, 195-211.
- Johnson M.E. & Hauser P.M. (2001). The practices of expert psychiatric nurses: Accompanying the patient to a calmer personal space. *Issues in Mental Health Nursing*, 22, 651-668.
- Izard, C. E., Fine, S., Mostow, A., Trentacosta, C., & Campbell, J. (2002). Emotion processes in normal and abnormal development and preventive intervention. *Development and Psychopathology*, 14, 761-787.
- Kempes, M., Matthys, W., deVries, H., van Engeland, H. (2005). Reactive and proactive aggression in children. A review of theory, findings, and the relevance for child and adolescent psychiatry. *European Journal of Child and Adolescent Psychiatry*, 14, 11-19.
- Lawson, L. (1998). Milieu management of traumatized youngsters. *Journal of Child and Adolescent Psychiatric nursing*, 11(3), 99-106
- Lazarus, R. S. & Folkman, S. (1984). *Stress, appraisal and coping*. New York: Springer.
- Lochman, J.E., Barry, T. D., & Pardini, D. A. (2003). Anger control training for aggressive youth. In A. E. Kazdin & J. R. Weisz (Eds.). *Evidence-based psychotherapies for children and adolescents* (pp 263-281). New York: Guilford Press.
- Lowe T. (1992). Characteristics of effective nursing interventions in the management of challenging behavior. *Journal of Advanced Nursing*, 17, 1226-1232.

- Maier, G. (1996). Managing threatening behavior. The role of talk down and talk up. *Journal of Psychosocial Nursing, 34*, 25-30.
- Martin K. (1995). Improving staff safety through an aggression management program. *Archives in Psychiatric Nursing, 9*, 211–215.
- Measham T.J. (1995). The acute management of aggressive behavior in hospitalized children and adolescents. *Canadian Journal of Psychiatry, 40*, 330–336.
- Morrison E.F. (1998). The culture of caregiving and aggression in psychiatric settings. *Archives of Psychiatric Nursing, 12*, 21–31.
- Morrison E.F. & Love C.C. (2003). An evaluation of four programs for the management of aggression in psychiatric settings. *Archives of Psychiatric Nursing, 27*, 146–155.
- Muris, P. Merckelbach, H. & Walczak, S. (2002). Aggression and threat perception abnormalities in children with learning and behavior problems. *Child Psychiatry and Human Development, 33*, 147-163.
- Perry, B.D., Pollard, R. A., Blakley, T. L., Baker, W. I, & Vigilante, D. (1995). Childhood trauma, the neurobiology of adaptation and “Use-dependant” development of the brain: How states become traits. *Infant Mental Health Journal, 16*, 271-291
- Posner, M. I., & Rothbart, M. K. (2000). Developing mechanisms of self regulation. *Development and Psychopathology, 12*, 427-441.
- Quay, H. C. (1997). Inhibition and attention deficit hyperactivity disorder. *Journal of Abnormal Child Psychology, 25*, 7-13.
- Rieder, C. & Cicchetti, D. (1989). Organizational perspective on cognitive control functioning and cognitive affective balance in maltreated children. *Developmental Psychology, 25*, 382-292
- Ryan E. P., Hart V.S., Messick D.L., Aaron J. & Burnette M. (2004). A prospective study of assault against staff by youths in a state psychiatric hospital. *Psychiatric Services, 55*, 665–670.
- Schneider, W. J., Cavell, T. A., & Hughes, J. N. (2003). A sense of containment: Potential moderator of the relation between parenting practices and children's externalizing behavior. *Development and Psychopathology, 15*, 95-117.
- Schore, A. N. (2001). The effects of early relational trauma on right brain development, affect regulation and infant mental health. *Infant Mental Health Journal, 22*, 201-269.

- Schwartz, D. (2000). Subtypes of victims and aggressors in children's peer groups. *Journal of Abnormal Child Psychology*, 28, 181-192.
- Shipman, K., Zeman, J., Penza, S., & Champion, K. (2000). Emotion management skills in sexually maltreated girls: A developmental psychopathology perspective. *Development and Psychopathology*, 12, 47-62.
- Sukhodolsky, D. G., Cardona, L., & Martin, A. (2005). Characterizing aggressive and noncompliant behaviors in a children's psychiatric inpatient settings. *Child Psychiatry and Human Development*, 36, 177-193.
- Shields, A., & Cicchetti, D. (1998). Reactive aggression among maltreated children: The contributions of attention and emotion dysregulation. *Journal of Clinical Child Psychology*, 27, 381-395.
- Thompson, R. A. (1994). Emotion regulation: A theme in search of a definition. In N. Fox (Ed.), *The development of emotion regulation: Biological and behavioral considerations. Monograph of the Society for Research in Child Development*, Vol 59, No.2-3, (pp. 25-52). Chicago: University of Chicago Press.
- Vermetten, E. & Bremner, J. D. (2002). Circuits and systems in stress: I Preclinical studies. *Depression and Anxiety*, 15, 126-147.
- Vitaro, F., Gendreau, P., Tremblay, R. E., & Oligny, P. (1998). Reactive and proactive aggression differentially predict later conduct problems. *Journal of Clinical Psychology and Psychiatry*, 39, 377-385.
- Vitiello, B., Behar, D., Hunt, J., Stoff, D. & Ricciuti, A. (1990). Subtyping aggression in children and adolescents. *Journal of Neuropsychiatry*, 2, 189-192.
- Vivona, J. M., Ecker, B., Halgin, R. P., Cates, D., Garrison, W. T., & Friedman, M. (1995). Self-and other-directed aggression in child and adolescent psychiatric inpatients. *Journal of the American Academy of Child and Adolescent Psychiatry*, 34, 434-444.
- Washchbusch, D. A., Willoughby, M. T., & Pelham, W. E. (1998). Criterion validity and the utility of reactive and proactive aggression: Comparisons to Attention Deficit Hyperactivity Disorder, Oppositional Defiant Disorder, Conduct Disorder and other measures of functioning. *Journal of Clinical Child Psychology*, 27, 396-405.

**GUIDELINES FOR THE ASSESSMENT OF SUICIDE RISK DURING CHILD
AND ADOLESCENT INPATIENT PSYCHIATRIC TREATMENT**

DESCRIPTION/OVERVIEW:

In the United States, suicide is the third leading cause of death in youth ages 15-19. While the youth suicide rate in the US is decreasing in most age segments, the youth suicide rate is steadily increasing worldwide. Nursing care of the suicidal child or adolescent at the primary treatment level includes identification and referral to the least restrictive setting for acute crisis management. Secondary treatment consists of ensuring the youth's safety needs throughout the entire crisis period until suicidal impulses are diminished. Tertiary treatment includes ongoing case management of the child or adolescent post-discharge in the community setting.

Many adults and caregivers do not believe that younger children can become suicidal. Youth suicide risk assessment is conducted in a thorough clinical interview using multiple modalities. Once admitted to the inpatient unit the task is to monitor risk and be aware of and respond to any changes in the level of risk. Youth self report of suicidal ideation and imminence of suicidal actions are accurate indicators of suicidality. Professionals need to be aware of this and other risk factors for suicide e.g., history of mood disorders, substance abuse, self-harming behaviors, family discord, childhood abuse history.

DESIRED OUTCOME:

The child or adolescent will be assessed by the nursing staff throughout treatment for suicidal risk. During the course of hospitalization, the child or adolescent will feel safe and gain control over suicidal impulses. The child or adolescent will benefit from a therapeutic treatment environment and learn age-appropriate coping skills. There will be a reduction in the intensity of the youth's emotional distress. Discharge planning will assure that the child and adolescent will have access to appropriate interventions during times of crisis and during follow-up care in the community.

A. ASSESSMENT

Principle of Care	Practice
The child/adolescent and family can expect to have suicide risk (i.e., absent, mild, moderate, severe, extreme) continuously assessed from the point of admission through discharge.	<p>The nurse, in collaboration with the child/adolescent and family, will: Assess the patient's suicide risk on a continuing basis according to the degree of depression and suicidal risk the child/adolescent exhibits.</p> <p>The nurse will consider using a standardized suicide risk assessment tool.</p> <p>Assess cultural norms around suicide.</p>

	<p>Assess spirituality in regards to suicide.</p> <p>Assess the patient for signs and symptoms of depression, including:</p> <ul style="list-style-type: none"> -Sleep disturbance, appetite loss or increase, increased agitation or irritability -Helplessness/hopelessness/powerlessness -Morbid preoccupation -Self-mutilating behaviors (SMB) -Social withdrawal -Lowered self-esteem -Decreased school performance -Sad, depressed vocalizations <p>Understand behaviors/factors that are considered dimensions of risk, these include:</p> <ul style="list-style-type: none"> -Being in a high-risk populations (i.e., psychiatric co-morbidity, history of physical or sexual abuse, disrupted/poor relationships with peers or family, sexual identity crisis) -Low GAF for last six months (severity of cumulative stressors) -History of prior suicide attempts -Lethality of past suicide attempt -Access to means -History of substance abuse -History of impulsive behavior - Suicidal ideation -Strength of intent to die, motivation - Family history of suicide
--	--

B. DIAGNOSIS

Assessment could result in any number of nursing diagnoses, including but not limited to:

Risk for violence: Directed towards self

Hopelessness related to

Powerlessness related to

Alteration in emotional integrity related to feelings of despair

Ineffective individual coping due to emotional pain, altered thought-processes, altered self-concept

Self-Mutilation related to

Potential for self-injury related to depression and despair

C. PLANNING

Principle of Care	Practice
<p>Child/adolescent and family can expect to participate in establishing goals for treatment.</p>	<p>The nurse will:</p> <p>Collaborate with the child/adolescent / family and treatment team to determine the goals of hospitalization.</p> <p>Document the ongoing process of planning around the child/adolescent/safety needs including the provision of safe milieu and the degree of required nursing supervision.</p>

D. IMPLEMENTATION

Principle of Care	Practice
<p>Child/adolescent and family can expect the nursing staff to provide on-going patient education and information related to suicide risk, diagnosis and treatment approaches.</p> <p>Child/adolescent and family can expect that the staff will implement all necessary procedures to assure a safe milieu.</p>	<p>The nurse will:</p> <p>Establish a relationship with patient that conveys sincere interest and caring and transfers the belief that help is possible.</p> <p>Discuss with child/adolescent ways to deal with suicidal ideations.</p> <p>Help child/adolescent strengthen existing coping skills.</p> <p>Provide group activities that allow patient to freely express their feelings and learn from others.</p> <p>Provide and promote education of child/adolescent and family regarding crisis intervention, diagnosis and medication.</p> <p>Assure patient is maintained at a suicide risk level commensurate with risk assessment.</p> <p>Follow institutional policy on monitoring patients on suicide risk or precautions.</p>

E. EVALUATION

Principle of Care	Practice
<p>Child/adolescent and family can expect ongoing responses to safety needs by nursing staff.</p>	<p>The nurse will:</p> <ul style="list-style-type: none"> Monitor and evaluate response to treatment on an ongoing basis. Alter treatment approaches in order to achieve goal of strengthening coping skills. Assist the patient and family to identify support systems within the school and community following discharge.

RESOURCES

- American Academy of Child & Adolescent Psychiatry. (2001). Practice parameters for the assessment and treatment of children and adolescents with suicidal behavior. *Journal of the American Academy of Child & Adolescent Psychiatry, 40*, 24Ss-51S.
- American Academy of Pediatrics. (1999). Waiting from harm: Deliberate self-harm and suicide in young people- A review of the literature. *Journal of Psychiatric and Mental Health Nursing, 6*, 91-100.
- Beautrais, A. L. (2003). Suicide and serious suicide attempts in youth: A multi-comparison group study. *American Journal of Psychiatry, 160*, 1093-1099
- Beckinsale, P., Martin, G., & Clark, S. (2001). Youth suicide in general practice. *Australian Family Physician, 30*(4), 391-394.
- Berman, A. J., Jobes, D., & Silverman, M. M. (2006). *Adolescent suicide: Assessment and intervention. (2nd Ed)*. Washington, D.C.: American Psychological Association.
- Bratter, T. E. (1997). *Guidelines for clinicians working with gifted, suicidal adolescents*. Retrieved February 1997, from <http://www.onlinepsych.com>
- Bridge, J. A., Barbe, R. P., & Brent, D. A. (2005). Recent trends in suicide among U.S. males, 1992-2001. *Psychiatric Services, 56*, 522.
- Bridge, J. A., Goldstein, T. R., & Brent, D. A. (2006). Adolescent suicide and suicidal behavior. *The Journal of Child Psychology and Psychiatry, 47*, 372-394
- Everall, R. D. (2000). Suicide: An attempted solution. *The Alberta Counsellor, 25*(2), 5-12.
- Gould, M.S., Greenberg, T., Velting, D. M., & Shaffer, D. (2003). Youth suicide risk and preventive interventions: A review of the past 10 years. *Journal of the American Academy of Child and Adolescent Psychiatry, 42*, 386-405.
- Health Education Consultants. Henkeleman, J. J. (2002). Evaluation of the suicidal child: Summary paper. Retrieved May 16, 2003 from <http://www.amhb.ab.ca/chmh/resources>.
- Hogarth, C. R. (1991). *Adolescent psychiatric nursing*. St. Louis, MO: C. V. Mosby.
- Joint Commission on Accreditation of Healthcare Organizations (2005). *CD-ROM: Reducing the risk of suicide*. Oakbrook Terrace, IL: Author.

- Maltsberger, J. T. (1988). *Determining suicidal risk*. Retrieved February 1997, from <http://www.onlinepsych.com>, [mentalhealth.com](http://www.mentalhealth.com). Internet Mental Health (<http://www.mental/health.com>.)
- Papolos, D., Hennen, J. & Cockerham, M. S. (2005). Factors associated with parent-report suicide threats by children and adolescents with community-diagnosed bipolar disorder. *Journal of Affective Disorders*, 86, 267-275.
- Pfeffer, C. R., Normadin, L., Kakuma, R. (1994). Suicidal children grow up: Suicidal and behavior and psychiatric disorders among relatives. *Journal of the Academy of Child and Adolescent Psychiatry*, 33,1087-1097.
- Pfeffer, C. R. (2001). Diagnosis of childhood and adolescent suicidal behavior. Unmet needs for suicide prevention. *Biological Psychiatry*, 49, 1055-1061.
- Thatcher, W. G., Reininger, B.M., & Drane, J.W. (2002). Using path analysis to examine adolescent suicide attempts, life satisfaction, and health risk behavior. *Journal of School Health* 72 (2), 70-71.
- Will, S. I. (1993). Depression/suicide. In M. K. Hass (Ed.), *The school nurse's source book of individualized health care plans (pp.211-212)*. South Branch, MN: Sunrise River Press.
- Windle, R.C.& Windle, M. (1997). An investigation of adolescent's substance use behaviors, depressed affect and suicidal behaviors. *Journal of Child psychology and Psychiatry*, 38, 921-929.
- Wunderlich, U., Bronisch, T., Wittchen, H-U, & Carter, R. (2001). Gender differences in adolescents and young adults with suicidal behavior. *Acta Psychiatrica Scandinavica*, 104, 332-229.

GUIDELINES FOR THE CARE OF CHILDREN AND ADOLESCENTS WITH A HISTORY OF MALTREATMENT DURING INPATIENT PSYCHIATRIC TREATMENT

DESCRIPTION/OVERVIEW:

The most recent data on child maltreatment indicate that in 2002, state child protective services received some 1.8 million referrals for possible maltreatment; approximately half of which, 900,000 children, were determined to be victims of abuse and neglect. Of these children about 60% were neglected, 20% physically abused and 10% sexually abused; children 0-3 years had the highest rate of victimization. Prevalence rates of maltreatment are primarily derived from adult accounts of their childhood experiences. In a large retrospective study of 13,500 HMO members, 43% of participants reported having been maltreated, either abused or witnessing abuse. Rates for different types of abuse vary. The prevalence rate for child sexual abuse is estimated to range from 16.8% for women and 7.9% for men. Recently one state report estimated that up to 84% of children hospitalized on inpatient psychiatric units had experienced some form of trauma. Thus inpatient units must take care to create milieus where children are not re-traumatized and afforded the opportunity to heal.

DESIRED OUTCOME

The child/adolescent will cope effectively with trauma-related symptoms. Discharge planning will include provision of services that will assist the child/adolescent in accomplishing developmental tasks and have an age-appropriate adjustment to social network. The child/adolescent will increase their ability to can identify emotions and express them in appropriate ways. If indicated the child/adolescent will establish a personal safety strategy. The child/adolescent will be future oriented in a positive way.

A. ASSESSMENT

Principle of Care	Practice
The child/adolescent can expect to have treatment needs assessed at the point of referral through the discharge phase of treatment.	The nurse, in partnership with the child/adolescent/family and members of the multidisciplinary treatment team will assess:
Assessment must include documentation of maltreatment and its aftermath.	If the child/adolescent wants to or is willing to share trauma history.
	Pre-trauma status including: age, developmental level, family structure, sociocultural factors, cognitive level, emotional issues and behavioral patterns.
	History of the maltreatment including occurrences, sexual activities, methods used and current status of perpetrator.

	<p>Resolution of the maltreatment including sanctions imposed on perpetrator.</p> <p>Indications for mandated reporting of ongoing maltreatment.</p> <p>Coping and defensive methods employed by the child/adolescent following abuse.</p> <p>Symptoms of general anxiety.</p> <p>Behavior patterns post-abuse indicative of trauma which may be integrated or characterized as predominantly anxious, avoidant, disorganized or aggressive.</p> <p>Triggers related to abuse that may increase youths' difficulties, i.e., closed doors, presence of man or women, time of year.</p> <p>What has been helpful in past when the child/adolescent has experienced trauma related affects.</p>
--	--

B. DIAGNOSIS

Assessment could result in any number of nursing diagnoses, including but not limited to

- Rape trauma syndrome related to rape/repeated trauma
- Coping, ineffective, individual related to.....
- Coping, ineffective, family, disabling, related to abuse by family member, denial by family
- Thought processes, alternation in, related to overwhelming anxiety
- Self-concept disturbance, related to abuse
- Sleep pattern disturbance, related to anxiety/fear/depression
- Social isolation related to failure of significant others to protect or rescue
- Violence, high risk for, self-directed (or directed at others) related to abuse
- Body image disturbance related to trauma
- Impaired social interaction related to.....
- Parenting, altered, related to abusive behavior, denial of abuse or guilt.

C. PLANNING

Principle of Care	Practice
<p>The child/adolescent/family can expect to participate collaboratively in their health care.</p>	<p>The nurse will collaborate with the child/adolescent and family and the multidisciplinary treatment team to:</p> <p>Determine short-term and long-term goals and interventions around processing the maltreatment that are realistic and empirically-based.</p> <p>Advocate for interventions that build on patient, child and community strengths.</p>

D. IMPLEMENTATION

Principle of Care	Practice
<p>The child/adolescent/family can expect:</p> <p>A psychological and physically safe milieu.</p> <p>The identification, recovery and strengthening of personal inner resources.</p> <p>When appropriate, support for the recall and processing of the traumatic experience.</p> <p>Education/information related to maltreatment and therapeutic treatment approaches.</p> <p>The child/adolescent/family can expect to receive anticipatory guidance.</p> <p>The child/adolescent with acute symptoms of trauma associated with the sexual abuse can expect support for the recall and processing of the traumatic experience.</p>	<p>The nurse will:</p> <p>Establish a safe, trusting relationship with the child/adolescent.</p> <p>Collaborate with the youth, treatment team and significant others to provide structure and maintain a safe environment.</p> <p>Use psychotherapeutic interventions that:</p> <ul style="list-style-type: none"> -Offer relief from negative feelings -Develop self-control -Develop emotional regulation -Enhance self-esteem -Foster understanding of personal reactions to maltreatment -Develop coping skills for upsetting thoughts, feelings and behavior. <p>Encourage use of previously identified and utilized coping skills.</p> <p>Provide education of the client and family/guardian regarding potential needs.</p>

	<p>For the child/adolescent with acute symptoms of trauma associated with abuse, the nurse will insure that the client has a circumscribed safe, accepting treatment environment in which the abusive traumatic experience can be discussed and the associated feelings freely expressed in order to:</p> <ul style="list-style-type: none"> -Normalize experiences and symptoms -Decrease emotional distress -Decrease avoidance -Develop a safety plan -Practice skills and generalize in-session symptom reduction to real-life situations. <p>Assure that the youth/family has continued access to support and treatment.</p>
--	--

E. EVALUATION

Principle of Care	Practice
<p>The child/adolescent/family can expect ongoing responses to progress in attaining desired outcomes.</p>	<p>The nurse will:</p> <p>Monitor or evaluate responses to treatment in an ongoing fashion.</p> <p>Collaboratively alter the treatment approaches in order to achieve the goal of optimum effective coping and healing.</p>

RESOURCES

- Burgess, A. W., Hartman, C. R. & Clements, P. T. (1995). Biology of memory and childhood trauma. *Journal of Psychosocial Nursing and Mental Health Services*, 33, 16-26.
- Brown, E. J., Albrecht, A., Munoz-Silva, D. M., McQuaide, J. & Silva, R. (2003). Treatment of children exposed to trauma: Research findings and therapeutic guidelines. In R. Silva (Ed.) *Posttraumatic stress disorder in children and adolescents handbook* (pp. 237-287). New York, NY: Norton Medical Publications.
- Charney, D. S., Deutch, A. Y., Southwick, S. M., & Krystal, J. H. (1995). Neural circuits and mechanisms of Post-Traumatic Stress Disorder. In M. J. Friedman, D. S. Charney, & A. Y. Deutch (Eds.), *Neurobiological and clinical consequences of stress: From normal adaptation to PTSD*, (pp. 271-287). Philadelphia: Lippincott-Raven.
- Cicchetti, D. & Toth, S. L. (1995). A developmental psychopathology perspective on child abuse and neglect. *Journal of the American Academy of Child and Adolescent Psychiatry*, 34, 541-565.
- Cloitre, M., Koenen, K. C., Cohen, L. R. & Han, H (2002). Skill training in affective and interpersonal regulation followed by exposure: A phase-based treatment for PTSD related to childhood abuse. *Journal of Consulting and Clinical Psychology*, 70, 1067-1074.
- Cohen, J. A. & Mannarino, A. P. (2000). Predictors of treatment outcome in sexually abused children. *Child Abuse and Neglect*, 24, 983-994.
- Diehl, A. S., & Prout, M. F. (2002). Effects of Posttraumatic stress disorder and child sexual abuse on self-efficacy development. *American Journal of Orthopsychiatry*, 72, 262-265.
- Edwards, V. J., Holden, G. W., Felitti, V. J., & Anda, R. F. (2003). Relationship between multiple forms of child maltreatment and adult mental health in community respondents: Results from adverse childhood experiences study. *American Psychiatric Association*, 160, 1453-1460.
- Everstine, D. S. & Everstine, L. (1993). *Sexual trauma in children and adolescents*. New York: Brunner/Mazel.
- Finzli, R., Ram, A., Shnit, D., Har-Even, D., Tyano, S., & Weizman, A. (2001). Depressive symptoms and suicidality in physically abused children. *American Journal of Orthopsychiatry*, 71, 98-107.

- Gil, E. (1996). *Treating abused adolescents*. New York: The Guilford Press.
- Gunner, M. R., & Cheatham, C. L. (2003). Brain and behavior interface: Stress and the developing brain. *Infant Mental Health Journal, 24*, 195-211.
- Hunter, M. (1991). *Abused boys: The neglected victims of sexual abuse*. New York: Fawcett Columbine Books.
- Kolko, D. J., Seleyo, J. & Brown, E. J. (1999). The treatment histories and service experiences of physically and sexually abuse families: Description, correspondence and correlates. *Child Abuse and Neglect, 23*, 459-476.
- LaBel, J. & Goldstein, R. (2004). Cents and sensibilities: What leaders should know. As Referenced in Regan, K. M., Curtin, C., & Vorderer, L. (2006). Paradigm shifts in inpatient psychiatric case of children: Approaching child-and-family centered care. *Journal of Child and Adolescent Psychiatric Nursing, 19*, 29-40.
- Lane, W. G., Dubowitz, H. & Harrington, D. (2002). Child sexual abuse evaluations: Adherence to recommendations source. *The Journal of Child Sexual Abuse, 11*, 1-18.
- Lau, A. S. & Weisz, J. R. (2003). Reported maltreatment among clinic-referred children: Implications for presenting problems, treatment attrition and long-term outcomes. *Journal of the American Academy of Child and Adolescent Psychiatry, 42*, 1327-1334.
- Lawson, L (1998). Milieu Management of Traumatized Youngsters. *Journal of Child and Adolescent Psychiatric Nursing, 11*, 99-106.
- Myers, J.E.B., Berliner, L., Briere, J., Hendrix, T. C., Jenny, C., & Reid, T. A. (2002). *The APSAC Handbook on Child Maltreatment (2nd Ed.)*. Thousand Oaks, CA: Sage.
- Morris, P. A. (1994). Superkids: Short-term group therapy for children with abusive backgrounds. *Journal of Child and Adolescent Psychiatric Nursing, 7*, 25-31.
- Orbach, Y., Hershkowitz, I., Lamb, M. E., Sternberg, K. J., Esplin, P. W. & Horowitz, D. (2000). Assessing the value of structured protocols for forensic interviews of alleged child abuse victims. *Child Abuse and Neglect, 24*, 733-752.
- Putnam, F. (2003). Ten-year research update review: Child sexual abuse. *Journal of the American Academy of Child and Adolescent Psychiatry, 42*, 269-278.
- Riddle, C. R. (1994). Development of an adolescent inpatient sexual abuse group: Application of Lewin's model of change. *Journal of Child and Adolescent Psychiatric Nursing, 7*, 17-24.

- Regan, K. M., Curtin, C., Vorderer, L. (2006). Paradigm shifts in inpatient psychiatric case of children: Approaching child-and-family centered care. *Journal of Child and Adolescent Psychiatric Nursing, 19*, 29-40.
- Rivard, J. C., McCorkle, D., Duncan, M. E., Pasquale, L. E., Bloom, S. L., & Abramovitz, R. (2004). Implementing a trauma recovery model for youths in residential treatment. *Child and Adolescent Social Work Journal, 21*, 529-550.
- Saywitz, K. J., Mannarino, A. P., Berliner, L., & Cohen, J. A. (2000). Treatment for sexually abused children and adolescents. *American Psychologist, 55*, 1040-1049.
- Teicher, M. H., Anderson, S.L., Polcari, A., Anderson, C. M., & Navalta, C. P. (2002). Developmental neurobiology of childhood stress and trauma. *Psychiatric Clinics of North America, 25*, 297-326.
- Tiernay, J. A. (2000). Post-traumatic stress disorder in children: Controversies and unresolved issues. *Journal of Child and Adolescent Psychiatric Nursing, 13*, 147-158.
- US Department of Health and Human Services (2004). *Summary Child Maltreatment 2002, Administration for Children and Families*. Retrieved on April 2004 from www.acf.dhhs.gov/programs/cb/publications/cm02/summary.htm.
- Vermetten, E. & Bremner, J. D. (2002). Circuits and systems in stress: I Preclinical studies. *Depression and Anxiety, 15*, 126-147.
- Waibel-Duncan, M. K. (2001). Medical fears following alleged child abuse. *Journal of Child and Adolescent Psychiatric Nursing, 14*, 179-185.
- Widom, C. S. (1999). Posttraumatic stress disorder in abused and neglected children grown up. *American Journal of Psychiatry, 156*, 1223-1229.

**PRINCIPLES OF TRANSCULTURAL CARE WITH SPECIAL EMPHASIS ON
APPLICATION TO INPATIENT PSYCHIATRIC HOSPITALIZATION FOR
CHILDREN AND ADOLESCENTS**

DESCRIPTION OVERVIEW:

Transcultural care is a dynamic process aimed at providing culturally sensitive and culturally congruent care that supports and promotes well-being in ethnically and culturally diverse individuals, families and communities. By practicing in a culturally competent manner, nurses can assist in minimizing client stressors and can incorporate cultural practices that support optimal coping behaviors. Cultural competence involves being aware of the other, having knowledge of the other, developing skills in working with the other, seeking encounters with the other and having a desire to learn about the culture of the other.

DESIRED OUTCOME:

The child / adolescent / family will be involved in and receive care / treatment that is sensitive to their cultural background and needs. The child / adolescent / family will leave the health care encounter feeling that their cultural values, beliefs and practices have been listened to, respected and wherever possible, maintained during the health care encounter. The health care provider will develop a repertoire of culturally competent skills that can be transferred and used as a template for each successive encounter with individuals from diverse cultures.

A. ASSESSMENT

Principle of Care	Practice
<p>The child / adolescent / family can expect:</p> <p>To receive an assessment of cultural needs on admission and continuously throughout the health care encounter.</p> <p>To be an active participant in communicating their mental health needs from their cultural perspective.</p> <p>Nursing staff to advocate for the inclusion of the child / adolescent / family in the development of the plan of care which will be sensitive to client needs and cultural practices.</p>	<p>The nurse, in partnership with the child / adolescent and family will assess for:</p> <p>Significant historical events from culture of origin that may affect health care behaviors.</p> <p>Language barriers or misunderstandings that may affect health care behaviors.</p> <p>Cultural views about health, mental health and mental illness.</p> <p>Cultural values, beliefs and health care practices that may affect health care behaviors.</p>

	<p>Child / adolescent / family view of the health care encounter, recommended treatment and relationship with service providers.</p> <p>Encourage the child / adolescent / family to verbalize questions or discomfort with health care recommendations.</p>
--	--

B. DIAGNOSIS

Assessment could result in any number of nursing diagnoses, including but not limited to:

- Impaired communication related to cultural and / or language differences
- Anxiety related to unmet needs
- Noncompliance related to client and provider relationship
- Noncompliance related to cultural dissonance
- Powerlessness related to health care environment
- Spiritual distress related to separation from religious / or cultural ties and practices
- At risk for violence related to cultural distress
- At risk for violence related to feelings of hopelessness or powerlessness
- Coping, ineffective individual related to....
- Coping, ineffective family related to...
- Social isolation or impaired social interaction related to cultural dissonance

C. PLANNING

Principle of Care	Practice
<p>The child / adolescent / family can expect to participate collaboratively in their health care.</p>	<p>In planning care, the nurse will promote:</p> <p>Respect and preservation of the client's cultural orientation.</p> <p>Accommodation to client culture within the treatment environment.</p> <p>Restructuring of care that innovatively meets the mental health needs of the individual and family without compromising their values.</p>

<p>The nurse in working with culturally diverse clients acquires the necessary knowledge about the health and mental health beliefs of the population.</p>	<p>The nurse will:</p> <p>Assess personal knowledge about the culture or ethnic group.</p> <p>Attend relevant seminars or workshops to learn more about the cultural or ethnic group.</p> <p>Seek cultural encounters with the cultural or ethnic group in order to enhance knowledge and understanding.</p>
--	--

D. IMPLEMENTATION

Principle of Care	Practice
<p>The child / adolescent / family can expect to receive on-going education related to diagnosis and various therapeutic treatment options.</p>	<p>The nurse will</p> <p>Listen to and ask questions of the patient to promote open and responsive communication.</p> <p>Provide and promote education of the child / adolescent / family in all aspects of mental health care.</p> <p>Assess youth's understanding of all information discussed.</p> <p>Utilize alternative methods of communication or resources in the event that the youth does not understand information conveyed.</p>

E. EVALUATION

Principle of Care	Practice
The child / adolescent / family can expect on-going and timely responses to stated cultural mental health care needs	<p>The nurse will</p> <p>Monitor and evaluate response to treatment in an on-going fashion.</p> <p>In collaboration with the youth and treatment team, revise interventions as necessary to achieve treatment goals while preserving cultural values of the child / adolescent / family.</p>

RESOURCES

- American Nurses Association (1997). *Position Statement: Cultural diversity in nursing practice*. Washington, DC: Author
- Campinha-Bacote, J. (2003). *Many faces: Addressing diversity in health care*. Online Journal of Issues in Nursing 8(1). Available: http://nursingworld.org/ojin/topic20_2htm.
- Campinha-Bacote, J. (1998). *The Process of Cultural Competence in the Delivery of Healthcare Services: A Culturally Competent Model of Care (3rd ed.)*. Cincinnati: Transcultural C.A.R.E. Associates.
- Department of Health and Human Services (2001). *Mental health: Culture, race, and ethnicity. A supplement to mental health: A report of the Surgeon General*. Washington, DC: Author.
- Kleinman, A. (1980). *Patients and healers in the context of culture. An exploration of the borderland between anthropology, medicine, and psychiatry*. Berkeley, CA: University of California Press.
- Kleinman, A. (1988). *The illness narratives. Suffering, healing and the human condition*. New York: Basic Books.
- Leininger, M. (Ed.) (1991). *Cultural care diversity and universality: A theory of nursing*. New York: National League for Nursing Press.
- Luna, I. (2002). Diversity issues in the delivery of healthcare. *Lippincott's Case Management* 7(4), 138-143.
- Mendyka, B. (2000). Exploring culture in nursing: A theory-driven practice. *Holistic Nursing Practice* 15(1), 32-41.

DISCHARGE PLANNING DURING BRIEF INPATIENT CHILD AND ADOLESCENT PSYCHIATRIC TREATMENT

DESCRIPTION/OVERVIEW:

Discharge planning will be based on established discharge criteria and on short-term and intermediate goals to be achieved by the child/adolescent prior to discharge from nursing care. These goals may serve as guidelines for development of long-term goals for the child/adolescent and family to work on after discharge. Discharge planning should begin almost upon admission. As treatment progresses the client and / or family members leaning needs should be assessed. They should have the opportunity to ask questions of the provider and may be given written information for later reference. They should be provided education about symptoms, medications, triggers, the disease process, anticipatory guidance and plans for intervening as well as facilitating early intervention before a crisis state evolves.

DESIRED OUTCOME:

The child/adolescent and family will be involved in discharge planning at the time of admission and throughout hospitalization. A discharge plan will be developed that matches the intensity of services with the intensity of the child/adolescent's needs. Discharge services will support family and child for optimal functioning in the least restrictive environment.

A. ASSESSMENT

Principle of Care	Practice
The child/adolescent/family can expect to undergo an assessment of discharge planning needs at the time of admission and throughout hospitalization.	<p>The nurse, in collaboration with the family and multidisciplinary treatment team will assure that discharge criteria includes assessment of:</p> <p>The patient's level of functioning and the family's ability to meet his/her safety needs.</p> <p>The child/adolescent and family's learning needs.</p> <p>The child/adolescent and family's attitude toward after-care; and the ability to follow through with after- treatment regime.</p> <p>Cultural variables, personal, and socioeconomic factors.</p>

	Educational needs including need for changes in educational services.
	Potential barriers to accessing services.

C. DIAGNOSES

Assessment could result in any number of nursing diagnoses, including but not limited to:

Noncompliance with treatment regimen
 Management of therapeutic regimen
 Knowledge deficit (specify).....

C. PLANNING

Principle of Care	Practice
<p>The child/adolescent/family can expect</p> <p>To have a discharge plan initiated at the time of admission.</p> <p>To have a discharge plan which reflects available support systems, cultural variables, personal and socioeconomic factors.</p> <p>The child/adolescent's discharge summary will reflect destination upon discharge, nursing interventions while in hospital, aftercare, prescriptions, and teaching.</p>	<p>The nurse will:</p> <p>Collaborate with the patient, family, outpatient providers (school where indicated) and multidisciplinary team to establish a discharge plan which will address immediate and future needs of the child/adolescent/family.</p> <p>Document the ongoing process of discharge planning.</p> <p>Write a discharge summary that reflects the child/adolescent's response to nursing interventions, teaching accomplished, goals, and destination upon discharge.</p>

D. IMPLEMENTATION

Principle of Care	Practice
<p>1. The child/adolescent and family can expect to receive sufficient explanation of the discharge plan such that they will verbalize understanding of the treatment recommendations.</p>	<p>The nurse in collaboration with the multidisciplinary team will:</p> <p>Initiate the discharge planning process beginning at admission.</p> <p>Discuss with the patient and family, anticipated learning needs related to discharge.</p> <p>Assure there has been adequate review of medications.</p>

	<p>Document the identified post discharge health care provider, the appropriate agency referrals and/or support groups.</p> <p>Share pertinent information with other disciplines and make recommendations regarding the youth's after care needs.</p>
--	--

E. EVALUATION

Principle of Care	Practice
The child/adolescent and family can expect to participate in the evaluation of the discharge plan.	The nurse will evaluate the appropriateness of the child/ adolescent's discharge plan based on the child/adolescent's needs, socioeconomic status, support system, educational/vocational abilities, and community resources.